
Terminology / Terminologie

Allergen nomenclature*

IUIS/WHO Allergen Nomenclature Subcommittee¹

The revised nomenclature for allergens is presented together with proposed nomenclatures for (a) allergen genes, mRNAs and cDNAs, and (b) recombinant and synthetic peptides of allergenic interest.

Introduction

Rapid advances have been made in the past few years on allergen characterization and sequence determination by chemical and molecular biological approaches. This is indicated by the list of allergens with known partial or complete amino acid sequences in Table 1. A number of other important allergens are known in addition to those in Table 1 but their sequences are as yet not known. A useful source for known allergens is the Allergen Database (ALBE) in which are compiled their known biochemical and immunological properties together with their sequence data if known (1).

To take into account these advances, a revision of the current allergen nomenclature system (2) is given below. As in the current nomenclature system, the proposed revisions are for allergens which induce IgE-mediated (atopic) allergy in humans. In addition to the expected thorough immunochemical characterization of any newly discovered allergen, investigators are urged to obtain partial, or preferably complete, sequence data before using the official nomenclature system. Also it is expected that investigators would screen a reasonable population size so as to establish the frequency of response in patients to the newly discovered allergens.

Investigators frequently refer to allergens as major or minor ones. The generally accepted mean-

ing of this terminology is that an allergen is designated as either major or minor depending on whether greater or less than 50% of patients tested have the corresponding allergen-specific IgEs (cf. 3-5).

The revised nomenclature for allergens is given below together with the proposed nomenclatures for (a) allergen genes, mRNAs and cDNAs and (b) recombinant and synthetic peptides of allergenic interest.

Revised nomenclature

(1) Allergens

Allergens are designated according to the accepted taxonomic name of their source as follows: the first three letters of the genus, space, the first letter of the species, space, and an Arabic number. The numbers are assigned to the allergens in the order of their identification, and the same number is generally used to designate homologous allergens of related species. As two examples, Lol p 1 refers to the first pollen allergen identified from *Lolium perenne*, rye grass, and Cyn d 1 refers to the homologous pollen allergen from *Cynodon dactylon*, Bermuda grass.

In some instances, the above system of the first 3 letters of a genus and the first letter of a species has to be modified to include an additional letter for designation of the exact genus or species. For example, 4 of the many vespids which can cause insect allergy are *Vespula vulgaris*, *Vespula vidua*, *Vespula consobrina* and *Vespa crabo*. The homologous major venom allergens, antigen 5s, from *Vespula vulgaris* and *Vespula vidua* are both designated as Ves v 5 in the existing nomenclature, and those from *Vespula consobrina* and *Vespa crabo* are designated as Ves c 5. To avoid these ambiguities, antigen 5s from *Vespula vulgaris* and *Vespula vidua* will be designated

* A résumé in French of this article appears on page 802. Requests for reprints should be addressed to the Chairman of the IUIS Nomenclature Committee, Professor Michel Kazatchkine, Unité d'Immunologie, Hôpital Broussais, 96 rue Didot, 75014 Paris, France.

¹ Members of this Nomenclature Subcommittee: Te Piao King (Chairman), Donald Hoffman, Henning Lowenstein, David G. Marsh, Thomas A.E. Platts-Mills and Wayne Thomas.

Reprint No. 5531

Table 1: Some allergens with known amino acid sequences

Allergen source	Allergens; systematic and original names	MW (kDa)	Sequence data ^{a, b}	References ^a
A. Weed pollens				
<i>Asterales</i>				
<i>Ambrosia artemisiifolia</i> (short ragweed)				
	Amb a 1; antigen E	38	C	8, 20
	Amb a 2; antigen K	38	C	8, 21
	Amb a 3; Ra3	11	C	22
	Amb a 5; Ra5	5	C	11, 23
	Amb a 6; Ra6	10	C	24, 25
	Amb a 7; Ra7	12	P	26
	Amb a ?	11	C	27
<i>Ambrosia trifida</i> (giant ragweed)				
	Amb t 5; Ra5G	4.4	C	9, 10, 28
<i>Artemisia vulgaris</i> (mugwort)				
	Art v 2	35	P	29
B. Grass pollens				
<i>Poales:</i>				
<i>Cynodon dactylon</i> (Bermuda grass)				
	Cyn d 1	32	C	30, 31
<i>Dactylis glomerata</i> (orchard grass)				
	Dac g 1; AgDg1	32	P	32
	Dac g 2	11	C	33
	Dac g 5		P	34
<i>Lolium perenne</i> (rye grass)				
	Lol p 1; group I	27	C	35, 36
	Lol p 2; group II	11	C	37
	Lol p 3; group III	11	C	38
	Lol p 5	31	P	34
	Lol p 9; Lol p 1b	31/35	C	39
<i>Phleum pratense</i> (timothy)				
	Phl p 1	27	C	40, 41
	Phl p 5; Ag25	32	C	42-45
<i>Poa pratensis</i> (Kentucky blue grass)				
	Poa p 1; group I	33	P	46
	Poa p 5	31	P	34
	Poa p 9	32/34	C	47
<i>Sorghum halepense</i> (Johnson grass)				
	Sor h 1		C	48
C. Tree pollens				
<i>Fagales:</i>				
<i>Alnus glutinosa</i> (alder)				
	Aln g 1	17	C	49
<i>Betula verrucosa</i> (birch)				
	Bet v 1	17	C	50
	Bet v 2; profilin	15	C	51
<i>Carpinus betulus</i> (hornbeam)				
	Car b 1	17	C	52
<i>Corylus avellana</i> (hazel)				
	Cor a 1	17	C	53
<i>Quercus alba</i> (white oak)				
	Que a 1	17	P	54
<i>Pinales:</i>				
<i>Cryptomeria japonica</i> (sugi)				
	Cry j 1	41-45	C	55, 56
	Cry j 2		C	57, 58
<i>Juniper sabinoides</i>				
	Jun s 1		C	59

(Table 1: continued)

<i>Juniper virginiana</i>	Jun v 1		C	58
Oleales:				
<i>Olea europea</i> (olive)	Ole e 1		C	59, 60
D. Mites				
<i>Dermatophagoides pteronyssinus</i> (mite)	Der p 1; antigen P1	25	C	61
	Der p 2	14	C	62
	Der p 3; trypsin	28/30	P	63
	Der p 4; amylase	60	P	64
	Der p 5	14	C	65
	Der p 6; chymotrypsin	25	P	66
	Der p 7	22-28	C	67
<i>Dermatophagoides microceras</i> (mite)	Der m 1	25	P	68
<i>Dermatophagoides farinae</i> (mite)	Der f 1	25	C	69
	Der f 2	14	C	70, 71
	Der f 3	30	P	72
<i>Lepidoglyphus destructor</i> (storage mite)	Lep d ?	15	P	73
E. Animals				
<i>Canis domesticus</i> ^c	Can d 1		C	74
	Can d 2		C	74
<i>Felis domesticus</i> (cat saliva)	Fel d 1; cat-1	38	C	15
<i>Mus musculus</i> (mouse urine)	Mus m 1; MUP	19	C	75, 76
<i>Rattus norvegicus</i> (rat urine)	Rat n 1	17	C	77, 78
F. Fungi				
<i>Aspergillus fumigatus</i>	Asp f 1	18	C	79
	Asp f ?	90	P	80
	Asp f ?	55	P	81
<i>Candida albicans</i>	Cand a ?	40	C	82
<i>Alternaria alternata</i>	Alt a 1	28	P	83-85
<i>Trichophyton tonsurans</i>	Tri t 1	30	P	86
G. Insects				
<i>Apis mellifera</i> (honey bee)	Api m 1; phospholipase A2	16	C	87
	Api m 2; hyaluronidase	44	C	88
	Api m 4; melittin	3	C	89
<i>Bombus pennsylvanicus</i> (bumble bee)	Bom p 1; phospholipase	16	P	90
	Bom p 4; protease		P	90
<i>Blattaria germanica</i> (cockroach)	Bla g 2	20	C	91
<i>Chironomus thummi thummi</i> (midges)	Chi t 1; hemoglobin	16	C	92
<i>Dolichovespula maculata</i> (white face hornet)	Dol m 1; phospholipase A1	35	C	93
	Dol m 2; hyaluronidase	44	C	94
	Dol m 5; antigen 5	23	C	95, 96
<i>Dolichovespula arenaria</i> (yellow hornet)	Dol a 5; antigen 5	23	C	97

UIIS/WHO Allergen Nomenclature Subcommittee

(Table 1: continued)

<i>Polistes annularis</i> (wasp)	Pol a 1; phospholipase A1	5	P	98
	Pol a 2; hyaluronidase	44	P	98
	Pol a 5; antigen 5	23	C	97
<i>Polistes exclamans</i> (wasp)	Pol e 1; phospholipase A1	34	P	100
	Pol e 5; antigen 5	23	C	97
<i>Polistes fuscatus</i> (wasp)	Pol f 5; antigen 5	23	C	99
<i>Polistes metricus</i> (wasp)	Pol m 5; antigen 5	23	P	99
<i>Vespula flavopilosa</i> (yellowjacket)	Ves f 5; antigen 5	23	C	99
<i>Vespula germanica</i> (yellowjacket)	Ves g 5; antigen 5	23	C	99
<i>Vespula maculifrons</i> (yellowjacket)	Ves m 1; phospholipase A1	33.5	C	101
	Ves m 2; hyaluronidase	44	P	102
	Ves m 5; antigen 5	23	C	97
<i>Vespula pennsylvanica</i> (yellowjacket)	Ves p 5; antigen 5	23	C	99
<i>Vespula squamosa</i> (yellowjacket)	Ves s 5; antigen 5	23	C	99
<i>Vespula vidua</i> (wasp)	Ves vi 5	23	C	99
<i>Vespula vulgaris</i> (yellowjacket)	Ves v 1; phospholipase A1	35	C	98
	Ves v 2; hyaluronidase	44	P	98
	Ves v 5; antigen 5	23	C	97
<i>Vespa crabo</i>	Vesp c 1; phospholipase	34	P	100
	Vesp c 5.0101; antigen 5	23	C	99
	Vesp c 5.0102; antigen 5	23	C	99
<i>Solenopsis invicta</i> (fire ant)	Sol i 2	13	C	103, 104
	Sol i 3	24	C	103
	Sol i 4	13	C	103
H. Foods				
<i>Gadus callarias</i> (cod)	Gad c 1; allergen M	12	C	105
<i>Gallus domesticus</i> (chicken)	Gal d 1; ovomucoid	28	C	106, 107
	Gal d 2; ovalbumin	44	C	106, 107
	Gal d 3; conalbumin (Ag22)	78	C	106, 107
	Gal d 4; lysozyme	14	C	106, 107
<i>Penaeus aztecus</i> (brown shrimp)	Pen a 1	36	P	108
	Pen a 2; tropomyosin	34	P	109
<i>Brassica juncea</i> (oriental mustard)	Bra j 1; 2S albumin	14	C	110
<i>Hordeum vulgare</i> (barley)	Hor v 1; BMAI-1	15	C	111
<i>Sinapis alba</i> (yellow mustard)	Sin a 1; 2S albumin	14	C	112
I. Others				
<i>Ascaris suum</i>	Asc s 1	10	P	113
<i>Hevea brasiliensis</i>	Hev b 1; elongation factor	58	P	114

^a References are those where partial (P) or complete (C) amino acid sequence data are available. Original references describing the initial characterization studies are not given because of limited space. We apologize to our colleagues whose allergen sequence data we may have overlooked for inclusion in this Table.

^b Sequence data for group 5 and 9 allergens from several grass pollens indicate that they are highly homologous proteins. Comparison of complete sequence data of group 5 and 9 allergens from a single grass species will clarify whether these two groups are the same protein.

^c *Canis domesticus* is also designated as *Canis familiaris*.

as Ves v 5 and Ves vi 5 respectively, and those from *Vespula consobrina* and *Vespa crabo* as Ves c 5 and Vesp c 5. In the examples given, the modified nomenclature is used for the allergens from *Vespula vidua* and *Vespa crabo*, as the allergens from *Vespula vulgaris* were characterized prior to those for *Vespula vidua* and *Vespa crabo*.

Another example is for allergens from the domestic dog (*Canis domesticus*) and the mold *Candida albicans*. To avoid ambiguity, the modified system is used to designate Can d and Cand a allergens from these two sources respectively.

In the current nomenclature system (2) the letters are italicized and the numerals are Roman numerals. In the revised system, only letters of normal type and Arabic numbers are used. The proposed changes conform to the accepted nomenclature used in bacterial genetics (6) and HLA system (7) in that italicized and normal characters are used to represent genotypes and phenotypes respectively.

(2) Allergens and isoallergens

An allergen from a single species may consist of several closely similar molecules. These similar molecules are designated as isoallergens when they share the following common biochemical properties: (a) similar molecular size; (b) identical biological function, if known, e.g. enzymatic action; and (c) >67% identity of amino acid sequences.

It is recognized that the recommended >67% sequence identity for 2 allergens to be assigned to the same group is only a guide. There are likely to be borderline cases. As an example, the ragweed allergens Amb a 1 and 2 share 65% amino acid sequence identity (8). These allergens were assigned to different groups because of their different immunochemical properties before their sequences were known.

Allergens from different species of the same or different genus which share the above-mentioned common biochemical properties are considered to belong to the same group and the sequence identity requirement can be <67% as is the case for allergens of the same species. For example, Amb a 5 and Amb t 5 from short and giant ragweed pollens have about 45% sequence identity (9), and also have similar tertiary structures (10, 11). Another example is the minor pollen allergens Amb a 10, Poa p 10 and Lol p 10 from ragweed (12), Kentucky bluegrass (13) and rye grass (14). Although their sequences are not known, they are assigned to the same allergen group as they clearly have the same biological function of cytochrome c.

(3) Isoallergens and variants

cDNA cloning of allergens often show nucleotide mutations which are either silent or which can lead

to single or multiple amino acid substitutions. In the revised system, members of an allergen group which have >67% amino acid sequence identity are designated as isoallergens. Each isoallergen may have multiple forms of closely similar sequences, which are designated as variants.

Isoallergens and their variants belonging to the same allergen group are designated by suffixes of a period followed by four Arabic numerals. The first two numerals 01 to 99 refer to a particular isoallergen, and the two subsequent numerals 01 to 99 refer to a particular variant of a particular isoallergen designated by the preceding two numerals. In cases where there is only one known isoallergen but there are several variants, the system of a suffix of 4 numerals will still apply. These numerals will be chosen in the order of the identification of allergens and/or their cDNAs irrespective of the physicochemical properties of the allergens. In cases of silent mutations, there can be more than one suffix of 4 numerals designating the same allergen and in that case the suffix with the lowest number will be used to designate the allergen of interest.

The addition of suffixes of 4 numerals to designate isoallergens and their variants will permit their unambiguous designation. In many cases it is unnecessary to specify the isoallergen or variant and the corresponding suffixes may be deleted; for example, Bet v 1 represents any Bet v 1 allergen and Bet v 1.0101 represents variant number 1 of isoallergen Bet v 1. Two other examples of this nomenclature system are given below.

On cDNA cloning, Amb a I showed multiple polymorphic forms which differ from each other by 12-24% in their sequences (8). Four such forms of Amb a I are known and they are designated as Amb a 1.01, 1.02, 1.03 and 1.04. Each isoallergen of the Amb a I group is found to have 1 to 3 variants with >97% of sequence identity. These variants of the Amb a 1.01 group will be designated as Amb a 1.0101, 1.0102 and etc.

In contrast to Amb a I, only two forms of Amb a II were found on cDNA cloning. These forms differ in two polymorphic sites and they have >99% amino acid sequence identity. They are designated as Amb a 2.0101 and 2.0102.

Nomenclature for allergen genes, mRNAs and cDNAs

At present the genomic structures of allergens are known in at least two cases; cat allergen Fel d 1 and mouse urinary allergen Mus m 1. Knowledge of the genomic structure can provide an understanding of how the different polymorphic forms are generated by differential splicing and/or exon usage. By adopt-

ing the revised nomenclature for allergens, we can reserve the italicized characters to designate an allergen gene. Normal characters are used for designation of mRNAs and cDNAs.

For example, Fel d 1 is a protein consisting of two polypeptide chains (15) which are encoded by two separate genes (16). The two allergen genes for chains 1 and 2 of Fel d 1 will be represented by *Fel d 1A* and *Fel d 1B* respectively. Allelic forms of mRNAs and cDNAs of the *Fel d 1A* gene are designated as mRNA or cDNA Fel d 1A.0101 where the numerals are to correspond to those of the polymorphic allergens.

Nomenclature for recombinant and synthetic peptides of allergenic interest

There is interest in the possible use of fragments of allergens as reagents to modulate allergen-specific immune responses. Such fragments may be prepared by recombinant technology or by chemical synthesis. Therefore it is useful to establish a generally accepted nomenclature for such peptides of allergenic interest.

The nomenclature for recombinant and synthetic peptides of allergenic interest is to be based on the nomenclature for naturally occurring allergens since it is well accepted by the scientific and the clinical communities. An allergen which is prepared by recombinant (r) or chemical synthetic (s) means is to be differentiated from a natural (n) allergen by the addition of the prefix of r or s followed by a suffix of the amino acid residue positions which are given in parenthesis. For examples, a recombinant hornet venom allergen Dol m 5.0201 which contains the entire sequence of 204 residues will be designated as rDol m 5.0201, and a recombinant or synthetic peptide of residue 151-165 of Dol m 5.0201 will be designated as r or sDol m 5.0201 (151-165) respectively.

Natural allergens may contain post-translational modifications as many proteins do. These modifications include glycosylation, acylation, methylation, etc. Recombinant or synthetic allergen, designated by the prefix r or s, is taken to indicate that it does not contain the post-translational modification of the natural allergen. If the recombinant or synthetic allergen does contain the exact same modification as that of the natural allergen, it will be designated with a prefix of rn or sn. For example, the honey bee venom allergen phospholipase A2, Api m 1, is a glycoprotein with an oligosaccharide attached at the asparagine residue number 13 (17, 18). A synthetic peptide of residues 1-20 of Api m 1 without the oligosaccharide at residue 13 will be designated as sApi m 1 (1-20), and a synthetic peptide of residue 1-20 with the exact same oligosaccharide of the natural allergen will be designated as snApi m 1 (1-20).

For recombinant or synthetic fragments which are derivatives of sequences contained within the native allergen structure, an additional suffix enclosed in square brackets will be used to indicate that the peptide referred to is an analogue. Substitutions or modifications of amino acid residues are given with the standard one-letter code and superscript numbers indicate the residue positions at which modifications occur. The one-letter codes for L-amino acids are given in upper case letters and those for D-amino acids are in lower case letters. The modifications which can be substitution, insertion or deletion are specified in parenthesis within the brackets. Obviously when there are many changes, it will not be practical to follow this nomenclature but to give the fully modified sequences. Examples of these analogues of sDol m 5.0201 (151-165) are given below:

Unmodified: sDol m 5.0201 (151-165)

Substitution: sDol m 5.0201 (151-165) [K¹⁵³]

— L-lysine residue at position 153 of sDol m 5.0201 (151-165) is substituted with D-lysine

Insertion: sDol m 5.0201 (151-165) [+K¹⁵³]

— one residue of L-lysine is inserted between positions 153-154

Deletion: sDol m 5.0201 (151-165) [-K¹⁵³]

— L-lysine residue at position 153 is deleted

N-terminal modification: sDol m 5.0201 (151-165) [N-Ac]

— N-terminal amino group is acetylated

C-terminal modification: sDol m 5.0201 (151-165) [C-NH₂]

— C-terminal carboxyl group is in the form of carboxamide

The nomenclature proposed above is very similar to that used for synthetic peptides representative of immunoglobulin sequences (19).

Acknowledgement

We thank Drs Martin Chapman, Irwin Griffith, Shayam Mohapatra, Rosalia Rodriguez and Alec Sehon for reviewing this article.

Résumé

Nomenclature des allergènes

La nomenclature des allergènes a été mise à jour en tenant compte des progrès rapides réalisés ces dernières années dans la caractérisation des allergènes et la détermination de leur séquence

en acides aminés par des techniques biochimiques et de biologie moléculaire.

La nouvelle nomenclature est présentée dans un tableau concernant les allergènes, les isoallergènes et leurs variants (pollens, acariens, autres allergènes, animaux, mycophytes, insectes, aliments, divers). La nomenclature concerne également a) les gènes, les ARNm et les ADNc des allergènes et b) les peptides synthétiques et recombinants importants en allergologie.

References

1. Marsh DG, Freidhoff LR. *ALBE, An allergen database*. IUIS, Baltimore, MD, Edition 1.0.
2. Marsh DG et al. Allergen nomenclature. *Bulletin of the World Health Organization*, 1986, **64**: 767-770.
3. King TP, PS Norman, Cornell JT. Isolation and characterization of allergen from ragweed pollen. II. *Biochemistry*, 1964, **3**: 458-468.
4. Lowenstein H. Timothy pollen allergens. *Allergy*, 1980, **35**: 188-191.
5. Aukrust L. Purification of allergens in *Cladosporium herbarum*. *Allergy*, 1980, **35**: 206-207.
6. Demerec M et al. A proposal for a uniform nomenclature in bacterial genetics. *Genetics*, 1966, **54**: 61-75.
7. Bodmer JG et al. Nomenclature for factors of the HLA system, 1990. *Immunogenetics*, 1991, **33**: 301-309.
8. Griffith IJ et al. Sequence polymorphism of Amb a I and Amb a II, the major allergens in *Ambrosia artemisiifolia* (short ragweed). *Int. arch. allergy appl. immunol.*, 1991, **96**: 296-304.
9. Roebber M et al. Immunochemical and genetic studies of Amb t V (Ra5G), an Ra5 homologue from giant ragweed pollen. *J. immunol.*, 1985, **134**: 3062-3069.
10. Metzler WJ et al. Solution structures of ragweed allergen Amb t V. *Biochemistry*, 1992, **31**: 5117-5127.
11. Metzler WJ et al. Proton resonance assignments and three-dimensional solution structure of the ragweed allergen Amb a V by nuclear magnetic resonance spectroscopy. *Biochemistry*, 1992, **31**: 8697-8705.
12. Goodfriend L et al. Cytochromes C: new ragweed pollen allergens. *Federation proceedings*, 1979, **38**: 1415.
13. Ekramoddoullah AKM, Kisil FT, Sehon AH. Allergenic cross reactivity of cytochrome c from Kentucky bluegrass and perennial ryegrass pollens. *Mol. immunol.*, 1982, **19**: 1527-1534.
14. Ansari AA, Killoran EA, Marsh DG. An investigation of human response to perennial ryegrass (*Lolium perenne*) pollen cytochrome c (Lol p X). *J. allergy clin. immunol.*, 1987, **80**: 229-235.
15. Morgenstern JP et al. Amino acid sequence of Fel d I, the major allergen of the domestic cat: protein sequence analysis and cDNA cloning. *Proc. Natl Acad. Sci. USA*, 1991, **88**: 9690-9694.
16. Griffith IJ et al. Expression and genomic structure of the genes encoding FdI, the major allergen from the domestic cat. *Gene*, 1992, **113**: 263-268.
17. Weber A, Marz L, Altmann F. Characteristics of the asparagine-linked oligosaccharide from honeybee venom phospholipase A2. *Comp. biochem. physiol.*, 1986, **83B**: 321-324.
18. Weber A et al. Specific interaction of IgE antibodies with a carbohydrate epitope of honey bee venom phospholipase A2. *Allergy*, 1987, **42**: 464-470.
19. Stanworth DR et al. Nomenclature for synthetic peptides representative of immunoglobulin chain sequences. *Bulletin of the World Health Organization*, 1990, **68**: 109-111.
20. Rafnar T et al. Cloning of Amb a I (Antigen E), the major allergen family of short ragweed pollen. *J. biol. chem.*, 1991, **266**: 1229-1236.
21. Rogers BL et al. Complete sequence of the allergen Amb a II: recombinant expression and reactivity with T cells from ragweed allergic patients. *J. immunol.*, 1991, **147**: 2547-2552.
22. Klapper DG, Goodfriend L, Capra JD. Amino acid sequence of ragweed allergen Ra3. *Biochemistry*, 1980, **19**: 5729-5734.
23. Ghosh B et al. Cloning and expression of immunologically active recombinant Amb a V allergen of short ragweed (*Ambrosia artemisiifolia*) pollen. *J. immunol.*, 1993, **150**: 5391-5399.
24. Roebber M et al. Isolation and properties of a new short ragweed pollen allergen, Ra6. *J. immunol.*, 1983, **131**: 706-711.
25. Lubahn B, Klapper DG. Cloning and characterization of ragweed allergen Amb a VI (abst). *J. allergy clin. immunol.*, 1993, **91**: 338.
26. Roebber M, Marsh DG. Isolation and characterization of allergen Amb a VII from short ragweed pollen. *J. allergy clin. immunol.*, 1991, **87**: 324.
27. Rogers BL et al. Cloning, complete sequence, and recombinant expression of a novel allergen from short ragweed pollen (abst). *J. allergy clin. immunol.*, 1993, **91**: 339.
28. Goodfriend L et al. Ra5G, a homologue of Ra5 in giant ragweed pollen: isolation, HLA-DR-associated activity and amino acid sequence. *Mol. immunol.*, 1985, **22**: 899-906.
29. Nilsen BM et al. Structural analysis of the glycoprotein allergen Art v II from pollen of mugwort (*Artemisia vulgaris*). *J. biol. chem.*, 1991, **266**: 2660-2668.
30. Matthiesen F, Schumacher M, Lowenstein H. Characterization of the major allergen of *Cynodon dactylon* (Bermuda grass) pollen. *J. allergy clin. immunol.*, 1991, **88**: 763-774.
31. Smith PM, Singh MB, Knox RB. Characterization and cloning of the major allergen of Bermuda grass, Cyn d I. In: Kraft D, Sehon A. eds. *Molecular biology and immunology of allergens*. Boca Raton, CRC Press, 1993: 157-160.
32. Mecheri S, Peltre G, David B. Purification and characterization of a major allergen from *Dactylis glomerata* pollen: the Ag Dg 1. *Int. archs allergy appl. immunol.*, 1985, **78**: 283-289.
33. Roberts AM et al. Nucleotide sequence of cDNA

- encoding the Group II allergen of Cocksfoot/Orchard grass (*Dactylis glomerata*), Dac g II. *European j. allergy clin. immunol.*, 1993, In press.
34. **Klynsner S et al.** Group V allergens in grass pollen IV. Similarities in amino acid compositions and amino terminal sequences of the group V allergens from *Lolium perenne*, *Poa pratensis* and *Dactylis glomerata*. *Clin. exp. allergy*, 1992, **22**: 491-497.
 35. **Perez M et al.** cDNA cloning and immunological characterization of the rye grass allergen Lol p I. *J. biol. chem.*, 1990, **265**: 16210-16215.
 36. **Griffith IJ et al.** Cloning and sequencing of Lol p I, the major allergenic protein of rye-grass pollen. *FEBS letters*, 1991, **279**: 210-215.
 37. **Ansari AA, Shenbagamurthi P, Marsh DG.** Complete amino acid sequence of a *Lolium perenne* (perennial rye grass) pollen allergen, Lol p II. *J. biol. chem.*, 1989, **264**: 11181-11185.
 38. **Ansari A.A, Shenbagamurthi P, Marsh DG.** Complete primary structure of a *Lolium perenne* (perennial rye grass) pollen allergen, Lol p III: Comparison with known Lol p I and II sequences. *Biochemistry*, 1989, **28**: 8665-8670.
 39. **Singh MB et al.** Isolation of cDNA encoding a newly identified major allergenic protein of rye-grass pollen: intracellular targeting to the amyloplast. *Proc. Natl Acad. Sci.*, 1991, **88**: 1384-1388.
 40. **Matthiesen F et al.** NH2-terminal sequences of four immunoaffinity purified grass pollen allergens: Phl p I, Poa p I, Sec c I and Cyn d I (abst.). *Allergy (Supp.)*, 1992, **47**: 31.
 41. **Laffer S et al.** cDNA cloning of a major allergen of rye (*Secale cereale*) and timothy grass (*Phleum pratense*). *XVth Eur. Congress of Allergology and Clin. Immunol.*, 1992, Paris.
 42. **Matthiesen F, Lowenstein H.** Group V allergens in grass pollens. I. Purification and characterization of the group V allergen from *Phleum pratense* pollen, Phl p V. *Clin. exp. allergy*, 1991, **21**: 297-307.
 43. **Vrtala S et al.** cDNA cloning of a major grass pollen allergen from timothy grass (*Phleum pratense*): identification as Phl p V, a protein with possible targeting to the amyloplast. *XVth Eur. Congress of Allergology and Clin. Immunol.*, 1992, Paris.
 44. **Peterson A, Becker WM, Schlaak M.** Characterization of isoforms of the major allergen Phl p V by immunoblotting and microsequencing. *Int. arch. allergy immunol.*, 1992, **98**: 105-109.
 45. **Bufe A et al.** Partial mRNA sequence of Phl p V. EMBL accession number X70942, 1993.
 46. **Esch RE, Klapper DG.** Isolation and characterization of a major cross-reactive grass group I allergenic determinant. *Mol. immunol.*, 1989, **26**: 557-561.
 47. **Olsen E et al.** Identification and characterization of the Poa p IX group of basic allergens of Kentucky bluegrass pollen. *J. immunol.*, 1991, **147**: 205-211.
 48. **Avjoglou A, Singh M, Knox RB.** Sequence analysis of Sor h I, the group I allergen of Johnson grass pollen and its comparison to rye-grass Lol p I (abst.). *J. allergy clin. immunol.*, 1993, **91**: 340.
 49. **Breiteneder H et al.** Complementary DNA cloning and expression in *Escherichia coli* of Aln g I, the major allergen in pollen of alder (*Alnus glutinosa*). *J. allergy clin. immunol.*, 1992, **90**: 909-917.
 50. **Breiteneder H et al.** The gene coding for the major birch pollen allergen Bet v I is highly homologous to a pea disease resistance response gene. *EMBO j.*, 1989, **8**: 1935-1938.
 51. **Valenta R et al.** Profilins constitute a novel family of functional plant pan-allergens. *J. exp. med.*, 1992, **175**: 377-385.
 52. **Larsen JN, Sturman P, Ipsen H.** PCR based cloning and sequencing of isogenes encoding the tree pollen major allergen Car b I from *Carpinus betulus*, hornbeam. *Mol. immunol.*, 1992, **29**: 703-711.
 53. **Breiteneder H et al.** Four recombinant isoforms of Cor a I, the major allergen of hazel pollen. *Europ. j. biochem.*, 1993, **212**: 355-362.
 54. **Ipsen H, Hansen BC.** The NH2-terminal amino acid sequence of the immunochemically partial identical major allergens of alder (*Alnus glutinosa*) Aln g I, birch (*Betula verrucosa*) Bet v I, hornbeam (*Carpinus betulus*) Car b I and oak (*Quercus alba*) Que a I pollens. *Molecular immunology*, 1991, **28**: 1279-1288.
 55. **Taniai M et al.** N-terminal amino acid sequence of a major allergen of Japanese cedar pollen (Cry j I). *FEBS letters*, 1988, **239**: 329-332.
 56. **Griffith IJ et al.** The cDNA cloning of Cry j I, the major allergen of *Cryptomeria japonica* (Japanese cedar) (abst.). *J. allergy clin. immunol.*, 1993, **91**: 339.
 57. **Sakaguchi M et al.** Identification of the second major allergen of Japanese cedar pollen. *Allergy*, 1990, **45**: 309-312.
 58. **Griffith I.** Personal communication.
 59. **Cardaba B et al.** Antibody response to olive pollen antigens: association between HLA class II genes and IgE response to Ole e I (abst.). *J. allergy clin. immunol.*, 1993, **91**: 338.
 60. **Villaalba M et al.** Amino acid sequence of Ole e I, the major allergen from olive tree pollen (*Olea europaea*). *Europ. j. biochem.* In press.
 61. **Chua KY, Stewart GA, Thomas WR.** Sequence analysis of cDNA encoding for a major house dust mite allergen, Der p I. *J. exp. med.*, 1988; **167**: 175-182.
 62. **Chua KY et al.** Isolation of cDNA coding for the major mite allergen Der p II by IgE plaque immunoassay. *Int. arch. allergy appl. immunol.*, 1990, **91**: 118-123.
 63. **Stewart GA et al.** The group III allergen from the house dust mite *Dermatophagoides pteronyssinus* is a trypsin-like enzyme. *Immunol.*, 1992, **75**: 29-35.
 64. **Lake FR et al.** House dust mite-derived amylase: allergenicity and physicochemical characterisation. *J. allergy clin. immunol.* 1991, **87**: 1035-1042.
 65. **Tovey ER et al.** Cloning and sequencing of a cDNA expressing a recombinant house dust mite protein that binds human IgE and corresponds to an important low molecular weight allergen. *J. exp. med.*, 1989, **170**: 1457-1462.
 66. **Yasueda H et al.** Allergenic and proteolytic properties of fourth allergens from *Dermatophagoides* mites. In: *Dust mite allergens and asthma. Report*

- of the 2nd International Workshop, Brussels, UCB Institute of Allergy, 1991: 63-64.
67. **Shen H-D et al.** Molecular cloning of a house dust mite allergen with common antibody binding specificities with multiple components in mite extracts. *Clin. exp. allergy*. In press.
 68. **Lind P, Hansen OC, Horn N.** The binding of mouse hybridoma and human IgE antibodies to the major fecal allergen, Der p I of *D. pteronyssinus*. *J. immunol.*, 1988, **140**: 4256-4262.
 69. **Dilworth RJ, Chua KY, Thomas WR.** Sequence analysis of cDNA coding for a major house dust allergen Der f I. *Clin. exp. allergy*, 1991, **21**: 25-32.
 70. **Nishiyama C et al.** Determination of three disulfide bonds in a major house dust mite allergen, Der f II. *Int. arch. allergy immunol.*, 1993, **101**: 159-166.
 71. **Trudinger M, Chua KY, Thomas WR.** cDNA encoding the major dust mite allergen Der f II. *Clin. exp. allergy*, 1991, **21**: 33-38.
 72. **Heymann et al.** Antigenic and structural analysis of group II allergens (Der f II and Der p II) from house dust mites (*Dermatophagoides spp.*). *J. allergy clin. immunol.*, 1989, **83**: 1055-1067.
 73. **van Hage-Hamsten M et al.** N-terminal amino acid sequence of major allergen of the mite *Lepidoglyphus destructor* (abst). *J. allergy clin. immunol.*, 1993, **91**: 353.
 74. **Konieczny A.** Personal communication.
 75. **McDonald B et al.** A 29 amino acid peptide derived from rat alpha 2 euglobulin triggers murine allergen specific human T cells (abst). *J. allergy clin. immunol.*, 1988, **83**: 251.
 76. **Clarke AJ et al.** Structure of mouse urinary protein genes: differential splicing configurations in the 3'-non-coding region. *EMBO j.*, 1984, **3**: 1045-1052.
 77. **Longbottom JL.** *Characterization of allergens from the urines of experimental animals*. London, Macmillan, 1983: 525-529.
 78. **Laperche Y et al.** Tissue-specific control of alpha 2u globulin gene expression: constitutive synthesis in submaxillary gland. *Cell*, 1983, **32**: 453-460.
 79. **Arruda LK, Mann BJ, Chapman MD.** Selective expression of a major allergen and cytotoxin, Asp f I in *Aspergillus fumigatus*. *J. immunol.*, 1992, **149**: 3554-3559.
 80. **Kumar A et al.** Isolation and characterization of a recombinant heat shock protein of *Aspergillus fumigatus*. *J. allergy clin. immunol.*, 1993, **91**: 1024-1030.
 81. **Teshima R et al.** Isolation and characterization of a major allergenic component (gp55) of *Aspergillus fumigatus*. *J. allergy clin. immunol.*, 1993, **92**: 698-706.
 82. **Shen HD et al.** The 40 kd allergen of *Candida albicans* is an alcohol dehydrogenase: molecular cloning and immunological analysis using monoclonal antibodies. *Clin. exp. allergy*, 1991, **21**: 675-681.
 83. **Matthiesen F, Olsen M, Lowenstein H.** Purification and partial sequence of the major allergen of *Alternaria alternata*, Alt a 1. *J. allergy clin. immunol.*, 1992, **89**: 241.
 84. **Curran IHA et al.** Purification and partial characterization of a low molecular weight antigen from *Alternaria alternata* (abst). *J. allergy clin. immunol.*, 1992, **89**: 283.
 85. **Vijay HM et al.** A major antigen of *Alternaria alternata* with potential for safe and effective immunotherapy. *J. allergy clin. immunol.*, 1993, **91**: 826-828.
 86. **Deuell B et al.** *Trichophyton tonsurans* allergen I. *J. immunol.*, 1991, **147**: 96-101.
 87. **Kuchler K et al.** Analysis of the cDNA for phospholipase A2 from honey bee venom glands: the deduced amino acid sequence reveals homology to the corresponding vertebrate enzymes. *Eur. j. biochem.*, 1989, **184**: 249-254.
 88. **Gmachl M, Kreil G.** Bee venom hyaluronidase is homologous to a membrane protein of mammalian sperm. *Proc. Natl Acad. Sci. USA*, 1993, **90**: 3569-3573.
 89. **Habermann E.** Bee and wasp venoms. *Science*, 1972, **177**: 314-322.
 90. **Jacobson RS, Hoffman DR.** Characterization of bumblebee venom allergens (abst). *J. allergy clin. immunol.*, 1993, **91**: 187.
 91. **Arruda LK, Vailes LD, Chapman MD.** Molecular cloning of cockroach (*B. germanica*) allergens (abst). *J. allergy clin. immunol.*, 1993, **91**: 188.
 92. **Mazur G, Baur X, Liebers V.** Hypersensitivity to hemoglobins of the Diptera family Chironomidae: structural and functional studies of their immunogenic/allergenic sites. *Monog. allergy*, 1990, **28**: 121-137.
 93. **Soldatova L, Kochoumian L, King TP.** Sequence similarity of a hornet (*D. maculata*) venom allergen phospholipase A1 with mammalian lipases. *FEBS letters*, 1993, **320**: 145-149.
 94. **Lu G, Kochoumian L, King TP.** White face hornet venom allergen, hyaluronidase: cloning and its sequence similarity with other proteins (abst.). 1994. *J. allergy clin. immunol.*, **93**: 224.
 95. **Fang KSF et al.** cDNA cloning and primary structure of a white-faced hornet venom allergen, antigen 5. *Proc. Natl Acad. Sci., USA*, 1988, **85**: 895-899.
 96. **King TP et al.** Structural studies of a hornet venom allergen antigen 5, Dol m V and its sequence similarity with other proteins. *Protein sequences and data analysis*, 1990, **3**: 263-266.
 97. **Lu G et al.** Sequence analysis and antigen cross reactivity of a venom allergen antigen 5 from hornets, wasps and yellowjackets. *J. immunol.*, 1993, **150**: 2823-2830.
 98. **King TP.** 1992. Unpublished data.
 99. **Hoffman DR.** Allergens in hymenoptera venom XXV: The amino acid sequences of antigen 5 molecules and the structural basis of antigenic cross-reactivity. *J. allergy clin. immunol.*, 1993, **92**: 707-716.
 100. **Hoffman DR.** 1992. Unpublished data.
 101. **Hoffman DR.** The complete amino acid sequence of a yellowjacket venom phospholipase (abst). *J. allergy clin. immunol.*, 1993, **91**: 187.
 102. **Jacobson RS, Hoffman DR, Kemeny DM.** The cross-reactivity between bee and vespid hyaluronidases has a structural basis (abst). *J. allergy clin. immunol.*, 1992, **89**: 292.
 103. **Hoffman DR.** Allergens in Hymenoptera venom

IUIS/WHO Allergen Nomenclature Subcommittee

- XXIV: The amino acid sequences of imported fire ant venom allergens Sol i II, Sol i III, and Sol i IV. *J. allergy clin. immunol.*, 1993, **91**: 71-78.
104. **Schmidt M et al.** Nucleotide sequence of cDNA encoding the fire ant venom protein Sol i II. *FEBS letters*, 1993, **319**: 138-140.
105. **Elsayed S et al.** The allergenic structure of allergen M from cod. Tryptic peptides of fragment TM1. *Int. arch. allergy appl. immunol.*, 1976, **52**: 59-63.
106. **Hoffman DR.** Immunochemical identification of the allergens in egg white. *J. allergy clin. immunol.*, 1983, **71**: 481-486.
107. **Langeland T.** A clinical and immunological study of allergy to hen's egg white. IV. specific IgE antibodies to individual allergens in hen's egg white related to clinical and immunological parameters in egg-allergic patients. *Allergy*, 1983, **38**: 493-500.
108. **Daul DB et al.** Common crustacea allergens: identification of B cell epitopes with the shrimp specific monoclonal antibodies. In: Kraft D, Sehon A, eds. *Molecular biology and immunology of allergens*. Boca Raton, CRC Press, 1993: 291-293.
109. **Rao PVS et al.** Tropomyosin is the major shrimp allergen (abst). *J. allergy clin. immunol.*, 1993, **91**: 341.
110. **Monsalve RI et al.** Characterization of a new mustard allergen, Bra j IE. Detection of an allergenic epitope. *Biochem. j.*, 1993, **293**: 625-632.
111. **Mena MR et al.** A major barley allergen associated with baker's asthma disease is a glycosylated monomeric inhibitor of insect α -amylase: cDNA cloning and chromosomal location of the gene. *Plant molec. biol.*, 1992, **20**: 451-458.
112. **Menendez-Arias L et al.** Primary structure of the major allergen of yellow mustard (*Sinapis alba* L.) seed, *Sin a I*. *Eur. j. biochem.*, 1988, **177**: 159-166.
113. **Christie JF et al.** N-terminal amino acid sequence identity between a major allergen of *Ascaris lumbricoides* and *Ascaris suum* and MHC-restricted IgE responses to it. *Immunology*, 1990, **69**: 596-602.
114. **Czuppon AB et al.** The rubber elongation factor of rubber trees (*Hevea brasiliensis*) is the major allergen in latex. *J. allergy clin. immunol.*, 1993, **92**: 690-697.